arXiv:1302.6650v1 [cond-mat.str-el] 27 Feb 2013

Anomalous Local Criticality in Heavy Fermion Metals from Holography

M.J.Luo[1]

[1] *Department of Physics, Tsinghua University, Beijing 100084, People's Republic of China*

We propose a holographic theory to explain numbers of anomalous critical phenomena observed in certain heavy-fermion metals, e.g. $\mathrm{CeCu_{5.9}Au_{0.1}}$ and $\mathrm{YbRh_2(Si_{0.95}Ge_{0.05})_2}$, incompatible with any conventional spin-density-wave quantum critical point theory. We show that the non-Gaussian nature of the fixed point from holography plays an essential role in the physics of these materials near a quantum critical point, which is not in the same universality class of the spin-density-wave type fixed point. The critical spin fluctuations at the non-Gaussian fixed point are strongly anisotropic, localized in spatial directions and critical in temporal directions with exponent $2/3$ in frequency over temperature dependence at low temperature. The local critical exponent tends unity which leads to a constant spin relaxation rate at high temperature quantum critical regime. The stability of the fixed point is also discussed.

Heavy fermion metals undergoing a magnetic quantum phase transition are expected well described by the Kondo lattice system of local moments/spins interacting with conducting electrons. However, certain heavy-fermion metals such as $\mathrm{CeCu_{6-x}Au_x}$ [1–4] and $\mathrm{YbRh_2(Si_{1-x}Ge_x)_2}$ [5–7] contradict the traditional Landau-Ginzburg type spin-density-wave (SDW) picture [8–12] (e.g. $\mathrm{CeNi_2Ge_2}$ [7], $\mathrm{CeCu_2Si_2}$ [13]) of magnetic quantum phase transition in the Kondo systems. Anomalous critical behaviors of spin susceptibility of these materials have been observed when they are chemically tuned to a quantum critical point (QCP) from paramagnetic metals to antiferromagnetism by varying x. A number of striking features shared by these materials near the QCP are inconsistent with usual SDW transition theory. (1) The frequency ω over temperature T scaling of the dynamical spin susceptibility displays a fractional exponent, however, ω/T scaling never occurs and its critical exponent is expected to take mean-field value of unity in 2D or 3D SDW QCP theories. (2) The fractional ω/T scaling exhibits wave vector independence, occurring not only at the antiferromagnetic wave vector but generic wave vector in the Brillouin zone. (3) Quasi-2D structure of these materials and strong anisotropic in critical spin fluctuation probed by neutron scattering experiment, which suggests certain local feature of the criticality. (4) Different from the traditional Kondo picture that the local spins of these materials are completely quenched by conducting electron at sufficient low temperature, in contrast, the spin susceptibility diverges rather saturates below the Kondo temperature which indicates the spins persist and a destruction of Kondo singlet near the QCP.

The feature (4) provides a clue to an important question that whether the spin critical point of these Kondo system at T=0 is asymptotic non-interacting one or interacting one. We can imagine that if the local spins in the Kondo system are asymptotically quenched in the vicinity of the QCP, the non-linear self-couplings between local spins are irrelevant in the renormalization-group sense, which naturally leads to a standard SDW type of QCP and hence this type of fixed point of the Kondo system is Gaussian. On the other hand, the fixed point could be non-Gaussian when the local spins are not free in the case of Kondo destruction. It is well known that different types of fixed point are associated with different universality classes taking different critical exponents, so the feature (1) and (4) are both imply the fixed point of these anomalous materials might not be trivially a Gaussian SDW type. The goal of this letter is to show that all these anomalous critical behaviors can be explained if we assume the Kondo system is at a non-Gaussian fixed point. However, studying the Kondo system at a non-Gaussian fixed point as a non-perturbative problem makes this idea difficult to be worked out theoretically. Phenomenologically, additional relevant critical modes responsible for the non-Gaussian aspects could be introduced based on certain effective models of the systems, the extended dynamical mean-field theory analysis of the Bose-Fermi Kondo model along this direction has been developed in [14–16], but so far to identify new critical modes in these systems experimentally is not an easy task. In this letter, we treat the Kondo lattice model as our first principle theory, and map the problem of Kondo systems at non-Gaussian fixed point to a solvable classical problems in a curved geometry from holographic duality.

We first generalize the spin symmetry of the Kondo system from SU(2) to SU(N) and take large-N [17], so in fact, the Kondo system is just a gauge system with SU(N) gauge group. The partitition function of Kondo system can be formally written as a functional of magnetic field coupled to the local spins by integrating out conduction electrons and local spins,

$$Z[B]=\int\prod_i\mathcal{D}[c^\dagger_{i\sigma},c_{i\sigma},S_{i\sigma\sigma'}]e^{-\beta\left[H[c^\dagger,c,S]+\mathrm{tr}(\mathbf{S}\cdot\mathbf{B})\right]},\tag{1}$$

in which H is the microscopic Kondo lattice Hamiltonian defined at ultraviolet,

$$H=\sum_{\langle ij\rangle\sigma}t_{ij}c^\dagger_{i\sigma}c_{j\sigma}+\frac{J_K}{N}\sum_{i\sigma\sigma'}\mathbf{S}_i\cdot c^\dagger_{i\sigma}\mathbf{\Gamma}_{\sigma\sigma'}c_{i\sigma'}+\frac{1}{N}\sum_{\langle ij\rangle}I_{ij}\mathbf{S}_i\cdot\mathbf{S}_j,\tag{2}$$

in which $c^{\dagger}_{i\sigma}, c_{i\sigma}$ are the conduction electrons at site i with spin index $\sigma = 1, ..., N$, S_i is the local spins at site i taken an antisymmetric representation of the SU(N) group, $\mathbf{\Gamma} = (\Gamma^1, \Gamma^2, ..., \Gamma^{N^2-1})$ are the SU(N) generator, J_k is the Kondo coupling between conduction electrons and local spins, I is the Ruderman-Kittel-Kasuya-Yosida (RKKY) coupling between nearest on site local spins, and B is the external magnetic field coupled with local spin, the trace is taken over the gauge representation of rank-N.

The details of the microscopic Hamiltonian may not be important, since the system becomes universal at fixed point including the infared non-Gaussian fixed point we focus in this letter, the important inputs are just the symmetry group and the dimensionality of the system. The holographic approach provides us a way to probe universal macroscopic features of such system at a non-Gaussian fixed point without knowing any microscopic details. The holographic duality says that there exists a non-Gaussian fixed point in this SU(N) gauge system at large-N limit, a d-dimensional theory at a vicinity of a non-Gaussian fixed point duals to a theory living on the boundary of an d+1 dimensional asymptotic anti-deSitter (AdS) gravitational background, $Z_{CFT} = Z_{AdS\,Gravity}$ [18, 19]. Therefore, the partition function $Z[B]$ at the non-Gaussian fixed point could be given by calculating its gravitational dual partition function. The large-N limit suppresses the fluctuations near the saddle point to the order $\mathcal{O}(1/N)$, the saddle point value gives the dominate contribution to the partition function of the gravity side, so the duality implies that the partition functional Eq.(1) takes the fixed point value

$$Z[B] = e^{-S_{cl}[B]}\Big|_{boundary}, \tag{3}$$

where S_{cl} is the classical action of magnetic field in an d+1 dimensional asymptotic AdS gravitational system. When the boundary theory becomes strongly coupled, the action is reduced to Einstein theory with classical asymptotic AdS metric. We take the U(1) subgroup of the SU(N), then

$$S_{cl}[B] = -\frac{1}{4g^2_{YM}}\int d^{d+1}x\sqrt{-g}F_{\mu\nu}F^{\mu\nu}, \tag{4}$$

where $F_{\mu\nu}$ is the fields strength, $B_i = \frac{1}{2}\epsilon_{ijk}F_{jk}$, $(i = 1, 2, 3)$, and $g^2_{YM} = 16\pi^2R/N^2$ is the gauge coupling. In order to model the quasi-2D structure and strong anisotropic mentioned in feature (3), in this letter we need $d = 3 + 1$ and lie the wave vector $\mathbf{q}$ along a special direction, e.g. $\mathbf{q}$ parallels to the easy moment direction c axis (we denote it as z-direction), $q^2 = |q_z|^2$. To introduce finite temperature to the system, a black hole is required to embed into the AdS_5 background, which gives the metric $g_{\mu\nu}$[20, 21]

$$ds^2 = \frac{(\pi TR)^2}{u}\left[-f(u)dt^2 + dx^2 + dy^2 + dz^2\right] + \frac{R^2}{4u^2f(u)}du^2, \tag{5}$$

where $f(u) = 1 - u^2$, and u valued from 0 (boundary) to 1 (horizon of the black hole) is the coordinate of the extra dimension, R is the curvature radius of the AdS space.

Thus far we are able to directly write down the non-Gaussian fixed point action of the Kondo system from the gravitation theory according to the holographic duality,

$$S_{cl}[B] = -\frac{N^2T^2}{16}\int dxdy\int duf(u)\int\frac{d\omega dq}{(2\pi)^2}\frac{1}{q^2}\left[B'^2_x(u,\tilde{\omega},\tilde{q}) + B'^2_y(u,\tilde{\omega},\tilde{q}) + ...\right], \tag{6}$$

where prime represents the derivative with respect to u, and $\tilde{\omega} = \omega/2\pi T$, $\tilde{q} = |q_z|/2\pi T$ are dimensionless frequency and wave vector scaled by temperature. Since the partition function is just simply a functional of B, we have formally integrated out all dynamic modes including fermionic and bosonic, then the effective spin susceptibility as the quadratic coefficient can be obtained by the linear-response theory from weak external applied magnetic field,

$$Z[B] = e^{-\int d^4x\langle S_i\rangle B_i + \frac{1}{2}\chi_{ij}B_iB_j + \mathcal{O}(B^3)...}, \tag{7}$$

where the Fourier components of the spin susceptibility χ is defined by the two-point function of spins,

$$\chi_{ij}(\omega,\mathbf{q}) = -i\int d\mathbf{x}dt\theta(t)\langle[S_i(t,\mathbf{x}), S_j(0,0)]\rangle e^{-i\omega t + i\mathbf{q}\cdot\mathbf{x}}. \tag{8}$$

Note that in Eq.(6) there are terms only with B_x and B_y, no B_z, we find

$$\chi = \chi_{xx} = \chi_{yy} = \frac{1}{q^2}\Phi(\tilde{\omega},\tilde{q})\Xi, \qquad \chi_{zz} = 0, \tag{9}$$

where

$$\Phi(\tilde{\omega}, \tilde{q}) = \lim_{u \to 0} \frac{B'_{x,y}}{B_{x,y}} \tag{10}$$

is a universal scaling function. This result Eq.(9) leads to an explanation of the feature (3), the strong anisotropic between $xx(yy)$ and zz components of the spin susceptibility is a direct consequence of the Bianchi identity or equivalently the transversality of the magnetic field propagating along z-direction. Do not confuse the anisotropy of the magnetic response with the isotropy of the AdS background Eq.(5), here the magnetic field in isotropic 3+1D reduces the critical spin system to an anisotropic XY spin system. $\Xi = \frac{1}{8}N^2T^2$ is the charge susceptibility [21]. Because of the sensitivity to the total number of degrees of freedom N and the non-trivial charge susceptibility $\Xi \sim \langle Q^2 \rangle$ at large-N and strong-coupling, the quantity we can use to compare with measurement is just $\chi_p = \chi/\Xi$.

The $\Phi(\tilde{\omega}, \tilde{q})$ is a regular function deduced from the solution of wave equation of the magnetic field, $\partial_\mu \left(\sqrt{-g} g^{\mu\nu} g^{\rho\lambda} F_{\nu\lambda}\right) = 0$, the wave equation in extra dimension is as

$$B''_{x,y} + \frac{f'}{f} B'_{x,y} + \frac{\tilde{\omega}^2 - \tilde{q}^2 f}{u f^2} B_{x,y} = 0. \tag{11}$$

As a consequence, $1/q^2$ is the only singular part of the χ_p in Eq.(9). It peaks when the wave vector along line in the specified z-direction taking critical wave vector, indicating the magnetic fluctuations and hence the spin fluctuations are quasi-2D (x-y plane) in real space. The singularity of χ_p in q-space with mean-field critical exponent signifies the emergence of spatially local critical modes that associated with the traditional long-wavelength fluctuations of the order parameter. The locally critical picture arise as a result of two-dimensional transverse magnetic fluctuations.

We are now in a position to consider the scaling function as a residue at the peak wave vector in Eq.(9), so we write it as $\Phi(\tilde{\omega}_q)$ in which we have taken the on-shell relation between ω and peaked q. Since the temperature is the only scale of the system, the $\Phi(\tilde{\omega}_q)$ exhibiting ω/T scaling is a direct consequence of conformal nature of the holographic theory in a thermo-bath. At low temperature limit, where $\tilde{\omega}, \tilde{q} \gg 1$, the magnetic field in bulk AdS space is relativistic, i.e. $\tilde{\omega}_q = \tilde{q}$. The asymptotic solution of Eq.(11) in this limit can be obtained by using Langer-Olver's method [22] [23], we find its spectral function

$$\mathrm{Im}\Phi(\tilde{\omega}_q) \propto \tilde{\omega}_q^{2/3}, \quad (\tilde{\omega}_q \gg 1). \tag{12}$$

The ω/T scaling exponent $2/3$ at the peak wave vector (on-shell modes) is an exact result in large-N limit corresponding to the case of strongly anisotropic XY spin system with completely vanished interplane response χ_{zz}. Such value of fractional scaling can also be obtained from numerical method. A deviation from the on-shell level condition has two important consequences, first, it leads to a correction to the on-shell value 2/3, i.e. $\mathrm{Im}\Phi \propto \tilde{\omega}^{2/3+\mathcal{O}(\tilde{\omega}-\tilde{q})}$; second, the exact transversality is also broken beyond the tree level which leads to a departure from the quasi-2D structure and strongly anisotropic, i.e. χ_{zz} deviates from zero.

Within experimental resolution, this on-shell value scaling agrees well with measurements of ac susceptibility $\chi_p \sim T^{-0.6}$ [6] in $\mathrm{YbRh_2(Si_{0.95}Ge_{0.05})_2}$, and approximated $\chi_p \sim T^{-0.75}$ [3] in $\mathrm{CeCu_{5.9}Au_{0.1}}$. The fractional exponent presented here demonstrates that it is not in the same universality class of the SDW theory nearby the antiferromagnetism. A salience feature of this result is that the anomalous exponent only occurs in the temporal fluctuations independent to the wave vector, while the exponent of spatial fluctuations are conventional mean-field type.

At high temperature limit, when $\tilde{\omega}, \tilde{q} \ll 1$, the dispersion is hydro-like, i.e. $\tilde{\omega}_q = \tilde{q}^2$, we find the scaling function

$$\mathrm{Im}\Phi(\tilde{\omega}_q) \propto \tilde{\omega}_q, \quad (\tilde{\omega}_q \ll 1). \tag{13}$$

Therefore, the scaling of local susceptibility gradually becomes unity in high temperature regime, which returns back to the standard Curie-Weiss scaling $\chi_p \sim T^{-1}$ and leads to a constant spin relaxation rate $T_1^{-1} \sim$ const.

Another important question concerns the stability of the non-Gaussian fixed point from holography. Considering modes propagating in an asymptotic AdS space with a charged black hole. The coupling between the modes and electric field of the charged black hole gives a contribution to the effective mass, $m_{eff}^2 = m^2 + e^2 g^{tt} \phi^2$, where ϕ is the electric field of the black hole, e the coupling constant. Note that the g^{tt} of AdS-charge black hole space is negative defined outside the horizon and diverges to $-\infty$ near the horizon, so the m_{eff}^2 is inevitable to become negative no matter how small the coupling and electric field are. Therefore, the modes are unstable even if the chemical potential that duals to the electric charge of the black hole is small, which leads to unstable modes living on the boundary of the AdS space. The instability is rather an intrinsic property of the holographic fixed point theory, so such fixed point we have discussed is in general associated with a real critical point observed in experiment but a phase (stable fixed point). In this sense, therefore, this type of quantum critical point is not very likely to develop superconducting phase close to the border of the antiferromagnetism, while the quantum critical point of the SDW type in a number of heavy-fermion metals do. Until now, no clearest experimental evidence shows that [24].

To summarize, we have generalized the Kondo lattice system to the SU(N) gauge theory with large-N, so beside a trivial Gaussian fixed point, a non-Gaussian fixed point of the system is also conjectured by the holographic duality. These two types of fixed points belong to different universality classes with different critical exponents. The non-Gaussian fixed point theory proposed by the holographic duality explains numbers of anomalous features observed in certain heavy-fermion metals (e.g. $\mathrm{CeCu_{5.9}Au_{0.1}}$ and $\mathrm{YbRh_2(Si_{0.95}Ge_{0.05})_2}$) that are incompatible with any SDW pictures. The feature (1): the ω/T scaling is the consequence of thermal conformal field nature of the holographic theory. The fractional critical exponent $\mathrm{Im}\chi_p(\tilde{\omega}_q) \sim \tilde{\omega}_q^{2/3}$ at peak wave vector, which fit the measurement well, is obtained by the asymptotic solution solving from the on-shell wave equation of magnetic field propagating in AdS space at low temperature. The feature (2): the mean-field value for the spatial critical exponent $\chi_p(q) \sim 1/q^2$ gives rise to the divergence of the correlation length in space and standard long-wavelength modes, which coexist with anomalous temporal local critical modes. The feature (3): the quasi-2D structure and strongly anisotropic spin susceptibility are the direct consequences of the special direction of the wave vector of the magnetic field and its transverse nature. The feature (4): the holographic theory is non-confining, so it only concerns with the Kondo-singlet-destroying quantum criticality in which the local spins persist to asymptotically low energies and become critical. Our theory also gives a standard Curie-Weiss scaling and constant spin relaxation rate at high temperature in quantum critical regime, which is consistent with measurements. Our theory also suggests that the fixed point corresponding to the anomalous criticality is unstable, so that it is bare and not very likely to develop superconducting phase near the quantum critical point.

This work has been supported in part by NSFC No.11205149.

---

[1] H. v. Löhneysen, T. Pietrus, G. Portisch, H. G. Schlager, A. Schröder, M. Sieck, and T. Trappmann. Non-fermi-liquid behavior in a heavy-fermion alloy at a magnetic instability. *Phys. Rev. Lett.*, 72:3262–3265, May 1994.

[2] A. Schröder, G. Aeppli, E. Bucher, R. Ramazashvili, and P. Coleman. Scaling of magnetic fluctuations near a quantum phase transition. *Phys. Rev. Lett.*, 80:5623–5626, Jun 1998.

[3] A. Schröder, G. Aeppli, R. Coldea, M. Adams, O. Stockert, H. v. Löhneysen, E. Bucher, R. Ramazashvili, and P. Coleman. Onset of antiferromagnetism in heavy-fermion metals. *Nature (London)* , 407:351–355, September 2000.

[4] O. Stockert, H. v. Löhneysen, A. Rosch, N. Pyka, and M. Loewenhaupt. Two-dimensional fluctuations at the quantum-critical point of $\mathrm{cecu}_{6-x}\mathrm{au}_x$. *Phys. Rev. Lett.*, 80:5627–5630, Jun 1998.

[5] O. Trovarelli, C. Geibel, S. Mederle, C. Langhammer, F. M. Grosche, P. Gegenwart, M. Lang, G. Sparn, and F. Steglich. $\mathrm{ybrh_2si_2}$: Pronounced non-fermi-liquid effects above a low-lying magnetic phase transition. *Phys. Rev. Lett.*, 85:626–629, Jul 2000.

[6] P. Gegenwart, J. Custers, Y. Tokiwa, C. Geibel, and F. Steglich. Ferromagnetic quantum critical fluctuations in $\mathrm{ybrh_2(si_{0.95}ge_{0.05})_2}$. *Phys. Rev. Lett.*, 94:076402, Feb 2005.

[7] R. Küchler, N. Oeschler, P. Gegenwart, T. Cichorek, K. Neumaier, O. Tegus, C. Geibel, J. A. Mydosh, F. Steglich, L. Zhu, and Q. Si. Divergence of the grüneisen ratio at quantum critical points in heavy fermion metals. *Phys. Rev. Lett.*, 91:066405, Aug 2003

[8] John A. Hertz. Quantum critical phenomena. *Phys. Rev. B*, 14:1165–1184, Aug 1976.

[9] A. J. Millis. Effect of a nonzero temperature on quantum critical points in itinerant fermion systems. *Phys. Rev. B*, 48:7183–7196, Sep 1993.

[10] Mucio A. Continentino. Universal behavior in heavy fermions. *Phys. Rev. B*, 47:11587–11590, May 1993.

[11] M. Lavagna and C. Pépin. Critical phenomena near the antiferromagnetic quantum critical point of heavy fermions. *Phys. Rev. B*, 62:6450–6457, Sep 2000.

[12] T. Moriya and T. Takimoto. Anomalous Properties around Magnetic Instability in Heavy Electron Systems. *Journal of the Physical Society of Japan*, 64:960, March 1995.

[13] H. Q. Yuan, F. M. Grosche, M. Deppe, C. Geibel, G. Sparn, and F. Steglich. Observation of Two Distinct Superconducting Phases in $\mathrm{CeCu_2Si_2}$. *Science*, 302:2104–2107, December 2003.

[14] Q. Si, S. Rabello, K. Ingersent, and L. Smith. Locally critical quantum phase transitions in strongly correlated metals. In *APS Meeting Abstracts*, page 16013, March 2001.

[15] Q. Si, J. L. Smith, and K. Ingersent. Quantum Critical Behavior in Kondo Systems. *International Journal of Modern Physics B*, 13:2331–2342, 1999.

[16] Qimiao Si, Silvio Rabello, Kevin Ingersent, and J. Lleweilun Smith. Local fluctuations in quantum critical metals. *Phys. Rev. B*, 68:115103, Sep 2003.

[17] P. Coleman, C. Pépin, and A. M. Tsvelik. Supersymmetric spin operators. *Phys. Rev. B*, 62:3852–3868, Aug 2000.

[18] Juan Martin Maldacena. The large N limit of superconformal field theories and supergravity. *Adv. Theor. Math. Phys.*, 2:231–252, 1998.

[19] Edward Witten. Anti-de Sitter space and holography. *Adv. Theor. Math. Phys.*, 2:253–291, 1998.

[20] Giuseppe Policastro, Dam T. Son, and Andrei O. Starinets. From AdS/CFT correspondence to hydrodynamics. *JHEP*, 09:043, 2002.

[21] Dam T. Son and Andrei O. Starinets. Minkowski space correlators in AdS / CFT correspondence: Recipe and applications. *JHEP*, 0209:042, 2002.

[22] F.W.J.Olver. The Asymptotic Solution of Linear Differential Equations of the Second Order for Large Values of a Parameter. *Royal Society of London Philosophical Transactions Series A*, 247,307 (1954).

[23] G.Policastro and A.Starinets. On the Absorption by Near-Extremal Black Branes. *Nucl.Phys.B*, 610,117 (2001).

[24] P. Gegenwart, Q. Si, and F. Steglich. Quantum criticality in heavy-fermion metals. *Nature Physics*, 4:186–197, March 2008.